



The American College of
Obstetricians and Gynecologists
WOMEN'S HEALTH CARE PHYSICIANS

COMMITTEE OPINION

Number 549 • January 2013

(Replaces Committee Opinion Number 315, September 2005)

Committee on Obstetric Practice

This document reflects emerging clinical and scientific advances as of the date issued and is subject to change. The information should not be construed as dictating an exclusive course of treatment or procedure to be followed.

Obesity in Pregnancy

ABSTRACT: In the United States, more than one third of women are obese, more than one half of pregnant women are overweight or obese, and 8% of reproductive-aged women are extremely obese, putting them at a greater risk of pregnancy complications. Therefore, preconception assessment and counseling are strongly encouraged for obese women and should include the provision of specific information concerning the maternal and fetal risks of obesity in pregnancy, as well as encouragement to undertake a weight-reduction program. At the initial prenatal visit, height and weight should be recorded for all women to allow calculation of body mass index (calculated as weight in kilograms divided by height in meters squared), and recommendations for appropriate weight gain should be reviewed at the initial visit and periodically throughout pregnancy. Nutrition consultation should be offered to all overweight or obese women, and they should be encouraged to follow an exercise program. Pregnant women who have undergone bariatric surgery should be evaluated for nutritional deficiencies and the need for vitamin supplementation when indicated. Obese patients undergoing cesarean delivery may require thromboprophylaxis with pneumatic compression devices and unfractionated heparin or low molecular weight heparin. For all obese patients, anesthesiology consultation early in labor should be considered, and consultation with weight-reduction specialists before attempting another pregnancy should be encouraged.

The prevalence of obesity in the United States has increased dramatically over the past 25 years. The recent National Health and Nutrition Examination Survey found that in the United States, more than one third of women are obese, more than one half of pregnant women are overweight or obese, and 8% of reproductive-aged women are extremely obese (1). The obesity problem is greatest among non-Hispanic black women (50%) compared with Mexican American women (45%) and non-Hispanic white women (33%) (1).

In 2009, the Institute of Medicine (IOM) published revised pregnancy weight gain guidelines that are based on prepregnancy body mass index (BMI) ranges recommended by the World Health Organization. These ranges are independent of age, parity, smoking history, race, and ethnic background (2, 3). The revised IOM recommendations define *normal weight* as a BMI of 18.5–24.9, *overweight* as a BMI of 25–29.9, and *obesity* as a BMI of 30 or greater. The IOM guidelines do not differentiate between Class I obesity (BMI=30–34.9), Class II obesity (BMI=35–39.9), and Class III obesity (BMI≥40) (2).

Pregnancy Complications

Overweight and obese women are at increased risk of several pregnancy complications, including gestational diabetes mellitus, hypertension, preeclampsia, cesarean delivery, and postpartum weight retention (4–8). Similarly, fetuses of pregnant women who are overweight or obese are at increased risk of prematurity, stillbirth, congenital anomalies, macrosomia with possible birth injury, and childhood obesity (9, 10). Additional concerns include potential intrapartum, operative, and postoperative complications and difficulties related to anesthesia management. Obese women are also less likely to initiate and sustain breastfeeding (11).

Maternal Complications

In a prospective multicenter study of more than 16,000 patients, a BMI of 30–39.9 was associated with an increased risk of gestational diabetes mellitus (odds ratio [OR], 2.6 and 4.0), gestational hypertension (OR, 2.5 and 3.2), preeclampsia (OR, 1.6 and 3.3), and fetal macrosomia (OR, 1.7 and 1.9), when compared with a BMI of

less than 30 (7). In this same study, the cesarean delivery rate was 20.7% for women with a BMI of 29.9 or less, 33.8% for women with a BMI of 30–34.9, and 47.4% for women with a BMI of 35–39.9. Other studies have consistently reported higher rates of preeclampsia, gestational diabetes mellitus, and cesarean delivery (particularly for arrest of labor) in obese women than in nonobese women (4–6).

At least three cohort studies suggest that obesity is an independent risk factor for spontaneous abortion among women who undergo infertility treatment (12–14). In recognition of this association, it is recommended that health care providers encourage obese women to lose weight before beginning infertility therapy. Data also link obesity with spontaneous abortion among women who conceive naturally (15).

Fetal Complications

When counseling obese women about potential pregnancy complications, it is important to inform them of the associated fetal risks, including prematurity, stillbirth, congenital abnormalities (eg, neural tube defects), macrosomia, and childhood and adolescent obesity. Some studies have reported a higher rate of premature delivery for obese women than for women of normal weight (4, 16). However, in a study of more than 2,900 obese women, prepregnancy obesity was associated with a lower rate of spontaneous preterm birth (17). A large Swedish cohort study reported a greater risk of antepartum stillbirth among obese patients than among women who had a BMI of less than 20 (16).

Obese pregnant women are more likely to give birth to an infant with congenital anomalies (9), and obesity also lowers detection rates of fetal anomalies during prenatal ultrasonography (18). Data establish that the risk of neural tube defects among obese pregnant women is double that of pregnant women of normal weight after correcting for diabetes as a potential confounding factor (19–21). The benefit of the administration of folic acid doses higher than 400 micrograms per day has not been studied in obese pregnant women without diabetes.

Multiple studies have shown that maternal obesity and excessive weight gain during pregnancy are associated with large-for-gestational-age infants (5, 22–24). Furthermore, these large-for-gestational-age infants are at increased risk of childhood and adolescent obesity (6, 25, 26). Although the diagnosis of fetal macrosomia is imprecise, prophylactic cesarean delivery may be considered for suspected fetal macrosomia with estimated fetal weights greater than 5,000 g in women without diabetes and greater than 4,500 g in women with diabetes (27).

Intrapartum Complications

It is important to discuss potential intrapartum complications with obese women, such as the challenges associated with anesthesia management and the increased risk of complicated and emergent cesarean delivery. Other

potential problems include difficulty estimating fetal weight (even with ultrasonography) and the inability to obtain interpretable external fetal heart rate and uterine contraction patterns.

Anesthesia Management

If an anesthesiology consultation was not obtained antepartum, it should be conducted early in labor to allow adequate time for the development of an anesthetic plan. The use of epidural or spinal anesthesia is recommended in the obese pregnant patient when anesthesia is needed or elected; however it may be technically difficult or impossible to administer this type of anesthesia because of obscured landmarks, difficult positioning, and excessive layers of adipose tissue. Alternatively, the use of general anesthesia in obese pregnant women also poses several challenges, including difficult endotracheal intubation due to excessive tissue and edema (28) and intraoperative respiratory events from failed or difficult intubation (29).

Cesarean Delivery

Operative and postoperative complications among obese pregnant women include increased rates of excessive blood loss, operative time greater than 2 hours, wound infection, and endometritis (30–32). Sleep apnea occurring in this group of women may further complicate anesthetic management and postoperative care (33).

Obese women who require cesarean delivery have an increased incidence of wound breakdown and infections (31, 34). Antimicrobial prophylaxis is recommended for all cesarean deliveries unless the patient is already receiving appropriate antibiotics (eg, for chorioamnionitis) (35). For obese women who require cesarean delivery, consideration should be given to using a higher dose of preoperative antibiotics for surgical prophylaxis (36). Attempts to decrease the incidence of wound breakdowns and infections that have been studied include closure of the subcutaneous layers and the placement of subcutaneous drains. Investigators have demonstrated that suture closure of the subcutaneous layer after cesarean delivery in obese patients may lead to a significant reduction in the incidence of postoperative wound disruption (37, 38). However, postoperative placement of subcutaneous draining systems has not consistently been shown to be of value in reducing postcesarean delivery morbidity (39, 40).

Because of an increased risk of venous thromboembolism, placement of pneumatic compression devices before cesarean delivery is recommended for all women not already receiving thromboprophylaxis (41). Obese patients, especially those who are hospitalized and immobile, may be at increased risk of thromboembolism. Individual risk assessment will lead some health care providers to plan thromboprophylaxis with pneumatic compression devices and unfractionated heparin or low molecular weight (LMW) heparin in such patients (42).

However, cesarean delivery in the emergency setting should not be delayed because of the timing necessary to implement thromboprophylaxis (41). Postpartum unfractionated heparin or LMW heparin prophylaxis has been recommended for patients thought to be at high risk of venous thromboembolism (43, 44); however, data are insufficient to determine whether the benefits of unfractionated heparin or LMW heparin prophylaxis in this group of patients outweigh the risks (45, 46).

Because of the increased likelihood of complicated and emergent cesarean delivery, extremely obese women may require specific resources, such as additional blood products, a large operating table, and extra personnel in the delivery room. Particular attention to the type and placement of the surgical incision is needed (ie, placing the incision above the panniculus adiposus) (34, 47). There are additional logistical challenges to monitoring labor and performing an emergent cesarean delivery in the extremely obese patient. Therefore, these patients should be counseled about these possible complications of an emergent cesarean delivery.

Bariatric Surgery Considerations

The number of obese reproductive-aged women undergoing bariatric surgery is increasing (48). Although maternal gastrointestinal complications, including obstruction and hemorrhage, can occur as a result of this procedure, the incidence is infrequent (49, 50). Furthermore, pregnancies after bariatric surgery are less likely to be complicated by gestational diabetes mellitus, hypertension, preeclampsia, and macrosomia than are pregnancies of obese women who have not undergone such surgery (51, 52).

Pregnant women who have undergone bariatric surgery should be evaluated for nutritional deficiencies and the need for vitamin supplementation when indicated because they are at increased risk of deficiencies in iron, vitamin B₁₂, folate, vitamin D, and calcium. Women with a gastric band should be monitored by their general surgeons during pregnancy because adjustment of the band may be necessary (53). Bariatric surgery is not an independent indication for cesarean delivery (54).

Counseling

Obese women are at increased risk of several pregnancy complications; therefore, preconception assessment and counseling are strongly encouraged. Obstetricians should provide education about the possible complications and should encourage obese patients to undertake a weight-reduction program, including diet, exercise, and behavior modification, before attempting pregnancy. Specific medical clearance may be indicated for some patients.

At the initial prenatal visit, height and weight should be recorded for all women to allow calculation of BMI, and recommendations for appropriate weight gain should be reviewed both at the initial visit and periodically throughout pregnancy (3). The 2009 IOM guidelines

recommend a total weight gain of 15–25 lb (6.8–11.3 kg) for overweight women (BMI=25–29.9) and 11–20 lb (5.0–9.1 kg) for all obese women (BMI ≥30). (For an online BMI calculator, see <http://www.nhlbisupport.com/bmi>). Nutrition consultation should be offered to all overweight or obese women, and they should be encouraged to follow an exercise program. Nutrition and exercise counseling should continue postpartum and before attempting another pregnancy. Because these patients are at increased risk of emergent cesarean delivery and anesthetic complications, anesthesiology consultation early in labor should be considered (22).

Recommendations

Recommendations for obese women who are pregnant or planning a pregnancy include the following:

- Preconception assessment and counseling are strongly encouraged and should include the provision of specific information concerning the maternal and fetal risks of obesity in pregnancy and encouragement to undertake a weight-reduction program.
- At the initial prenatal visit, height and weight should be recorded for all women to allow calculation of BMI, and recommendations for appropriate weight gain, guided by IOM recommendations, should be reviewed both at the initial visit and periodically throughout pregnancy.
- Nutrition consultation should be offered to all overweight or obese women, and they should be encouraged to follow an exercise program. Nutrition and exercise counseling should continue postpartum and before attempting another pregnancy.
- Women who have undergone bariatric surgery should be evaluated for nutritional deficiencies and the need for vitamin supplementation, when indicated, because they are at increased risk of deficiencies in iron, vitamin B₁₂, folate, vitamin D, and calcium.
- For patients undergoing cesarean delivery who have additional risk factors for thromboembolism such as obesity, individual risk assessment may require thromboprophylaxis with pneumatic compression devices and unfractionated heparin or LMW heparin.
- Consideration should be given to using a higher dose of preoperative antibiotics for cesarean delivery prophylaxis.
- The use of suture closure of the subcutaneous layer after cesarean delivery in obese patients may lead to a significant reduction in the incidence of postoperative wound disruption.
- Anesthesiology consultation early in labor should be considered.
- Consultation with a weight-reduction specialist before attempting another pregnancy should be encouraged.

References

1. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA* 2012; 307:491–7. [PubMed] [Full Text] ↩
2. Institute of Medicine. Weight gain during pregnancy: reexamining the guidelines. Washington, DC: National Academies Press; 2009. ↩
3. Weight gain during pregnancy. Committee Opinion No. 548. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2013;121:210–2. [Obstetrics & Gynecology] ↩
4. Baeten JM, Bukusi EA, Lambe M. Pregnancy complications and outcomes among overweight and obese nulliparous women. *Am J Public Health* 2001;91:436–40. [PubMed] [Full Text] ↩
5. Cedergren MI. Maternal morbid obesity and the risk of adverse pregnancy outcome. *Obstet Gynecol* 2004;103:219–24. [PubMed] [Obstetrics & Gynecology] ↩
6. Sebire NJ, Jolly M, Harris JP, Wadsworth J, Joffe M, Beard RW, et al. Maternal obesity and pregnancy outcome: a study of 287,213 pregnancies in London. *Int J Obes Relat Metab Disord* 2001;25:1175–82. [PubMed] [Full Text] ↩
7. Weiss JL, Malone FD, Emig D, Ball RH, Nyberg DA, Comstock CH, et al. Obesity, obstetric complications and cesarean delivery rate—a population-based screening study. FASTER Research Consortium. *Am J Obstet Gynecol* 2004;190:1091–7. [PubMed] [Full Text] ↩
8. Vesco KK, Dietz PM, Rizzo J, Stevens VJ, Perrin NA, Bachman DJ, et al. Excessive gestational weight gain and postpartum weight retention among obese women. *Obstet Gynecol* 2009;114:1069–75. [PubMed] [Obstetrics & Gynecology] ↩
9. Stothard KJ, Tennant PW, Bell R, Rankin J. Maternal overweight and obesity and the risk of congenital anomalies: a systematic review and meta-analysis. *JAMA* 2009;301:636–50. [PubMed] [Full Text] ↩
10. Oken E, Taveras EM, Kleinman KP, Rich-Edwards JW, Gillman MW. Gestational weight gain and child adiposity at age 3 years. *Am J Obstet Gynecol* 2007;196:322.e1–8. [PubMed] [Full Text] ↩
11. Li R, Jewell S, Grummer-Strawn L. Maternal obesity and breast-feeding practices. *Am J Clin Nutr* 2003;77:931–6. [PubMed] [Full Text] ↩
12. Bellver J, Rossal LP, Bosch E, Zuniga A, Corona JT, Melendez F, et al. Obesity and the risk of spontaneous abortion after oocyte donation. *Fertil Steril* 2003;79:1136–40. [PubMed] [Full Text] ↩
13. Fedorcsak P, Storeng R, Dale PO, Tanbo T, Abyholm T. Obesity is a risk factor for early pregnancy loss after IVF or ICSI. *Acta Obstet Gynecol Scand* 2000;79:43–8. [PubMed] [Full Text] ↩
14. Wang JX, Davies MJ, Norman RJ. Obesity increases the risk of spontaneous abortion during infertility treatment. *Obes Res* 2002;10:551–4. [PubMed] ↩
15. Lashen H, Fear K, Sturdee DW. Obesity is associated with increased risk of first trimester and recurrent miscarriage: matched case-control study. *Hum Reprod* 2004;19:1644–6. [PubMed] [Full Text] ↩
16. Cnattingius S, Bergstrom R, Lipworth L, Kramer MS. Pre-pregnancy weight and the risk of adverse pregnancy outcomes. *N Engl J Med* 1998;338:147–52. [PubMed] [Full Text] ↩
17. Hendler I, Goldenberg RL, Mercer BM, Iams JD, Meis PJ, Moawad AH, et al. The Preterm Prediction Study: association between maternal body mass index and spontaneous and indicated preterm birth. *Am J Obstet Gynecol* 2005;192:882–6. [PubMed] [Full Text] ↩
18. Dashe JS, McIntire DD, Twickler DM. Effect of maternal obesity on the ultrasound detection of anomalous fetuses. *Obstet Gynecol* 2009;113:1001–7. [PubMed] [Obstetrics & Gynecology] ↩
19. Shaw GM, Velie EM, Schaffer D. Risk of neural tube defect-affected pregnancies among obese women. *JAMA* 1996;275:1093–6. [PubMed] ↩
20. Waller DK, Mills JL, Simpson JL, Cunningham GC, Conley MR, Lassman MR, et al. Are obese women at higher risk for producing malformed offspring? *Am J Obstet Gynecol* 1994;170:541–8. [PubMed] ↩
21. Werler MM, Louik C, Shapiro S, Mitchell AA. Prepregnant weight in relation to risk of neural tube defects. *JAMA* 1996;275:1089–92. [PubMed] ↩
22. Rode L, Nilas L, Wojdemann K, Tabor A. Obesity-related complications in Danish single cephalic term pregnancies. *Obstet Gynecol* 2005;105:537–42. [PubMed] [Obstetrics & Gynecology] ↩
23. Stephansson O, Dickman PW, Johansson A, Cnattingius S. Maternal weight, pregnancy weight gain, and the risk of antepartum stillbirth. *Am J Obstet Gynecol* 2001;184:463–9. [PubMed] [Full Text] ↩
24. Watkins ML, Rasmussen SA, Honein MA, Botto LD, Moore CA. Maternal obesity and risk for birth defects. *Pediatrics* 2003;111:1152–8. [PubMed] [Full Text] ↩
25. Hediger ML, Overpeck MD, McGlynn A, Kuczumarski RJ, Maurer KR, Davis WW. Growth and fatness at three to six years of age of children born small- or large-for-gestational age. *Pediatrics* 1999;104:e33. [PubMed] [Full Text] ↩
26. Salsberry PJ, Reagan PB. Taking the long view: the prenatal environment and early adolescent overweight. *Res Nurs Health* 2007;30:297–307. [PubMed] ↩
27. American College of Obstetricians and Gynecologists. Fetal macrosomia. ACOG Practice Bulletin 22. Washington, DC: ACOG; 2000. ↩
28. Mhyre JM. Anesthetic management for the morbidly obese pregnant woman. *Int Anesthesiol Clin* 2007;45:51–70. [PubMed] ↩
29. Hood DD, Dewan DM. Anesthetic and obstetric outcome in morbidly obese parturients. *Anesthesiology* 1993;79:1210–8. [PubMed] ↩
30. Kabiru W, Raynor BD. Obstetric outcomes associated with increase in BMI category during pregnancy. *Am J Obstet Gynecol* 2004;191:928–32. [PubMed] [Full Text] ↩
31. Myles TD, Gooch J, Santolaya J. Obesity as an independent risk factor for infectious morbidity in patients who undergo cesarean delivery. *Obstet Gynecol* 2002;100:959–64. [PubMed] [Obstetrics & Gynecology] ↩
32. Perlow JH, Morgan MA. Massive maternal obesity and peri-operative cesarean morbidity. *Am J Obstet Gynecol* 1994;170:560–5. [PubMed] ↩

33. Maasilta P, Bachour A, Teramo K, Polo O, Laitinen LA. Sleep-related disordered breathing during pregnancy in obese women. *Chest* 2001;120:1448–54. [PubMed] ↩
34. Wall PD, Deucy EE, Glantz JC, Pressman EK. Vertical skin incisions and wound complications in the obese parturient. *Obstet Gynecol* 2003;102:952–6. [PubMed] [Obstetrics & Gynecology] ↩
35. Antimicrobial prophylaxis for cesarean delivery: timing of administration. Committee Opinion No. 465. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2010;116:791–2. [PubMed] [Obstetrics & Gynecology] ↩
36. Use of prophylactic antibiotics in labor and delivery. Practice Bulletin No. 120. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2011;117:1472–83. [PubMed] [Obstetrics & Gynecology] ↩
37. Cetin A, Cetin M. Superficial wound disruption after cesarean delivery: effect of the depth and closure of subcutaneous tissue. *Int J Gynaecol Obstet* 1997;57:17–21. [PubMed] [Full Text] ↩
38. Chelmos D, Rodriguez EJ, Sabatini MM. Suture closure of subcutaneous fat and wound disruption after cesarean delivery: a meta-analysis. *Obstet Gynecol* 2004;103:974–80. [PubMed] [Obstetrics & Gynecology] ↩
39. Al-Inany H, Youssef G, Abd ElMaguid A, Abdel Hamid M, Naguib A. Value of subcutaneous drainage system in obese females undergoing cesarean section using Pfannenstiel incision. *Gynecol Obstet Invest* 2002;53:75–8. [PubMed] ↩
40. Magann EF, Chauhan SP, Rodts-Palenik S, Bufkin L, Martin JN Jr, Morrison JC. Subcutaneous stitch closure versus subcutaneous drain to prevent wound disruption after cesarean delivery: a randomized clinical trial. *Am J Obstet Gynecol* 2002;186:1119–23. [PubMed] [Full Text] ↩
41. Thromboembolism in pregnancy. Practice Bulletin No. 123. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2011;118:718–29. [PubMed] [Obstetrics & Gynecology] ↩
42. Bates SM, Greer IA, Hirsh J, Ginsberg JS. Use of anti-thrombotic agents during pregnancy: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004;126:627S–44S. [PubMed] ↩
43. Bates SM, Greer IA, Pabinger I, Sofaer S, Hirsh J. Venous thromboembolism, thrombophilia, antithrombotic therapy, and pregnancy: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest* 2008;133:844S–86S. [PubMed] [Full Text] ↩
44. Royal College of Obstetricians and Gynaecologists. Report of the RCOG working party on prophylaxis against thromboembolism in gynaecology and obstetrics. London (UK): RCOG; 1995. ↩
45. Tooher R, Gates S, Dowswell T, Davis LJ. Prophylaxis for venous thromboembolic disease in pregnancy and the early postnatal period. *Cochrane Database of Systematic Reviews* 2010, Issue 5. Art. No.: CD001689. DOI: 10.1002/14651858.CD001689.pub2. [PubMed] [Full Text] ↩
46. Hague WM, North RA, Gallus AS, Walters BN, Orlikowski C, Burrows RF, et al. Anticoagulation in pregnancy and the puerperium. Working Group of the Obstetric Medicine Group of Australasia. *Med J Aust* 2001;175:258–63. [PubMed] [Full Text] ↩
47. Houston MC, Raynor BD. Postoperative morbidity in the morbidly obese parturient woman: supraumbilical and low transverse abdominal approaches. *Am J Obstet Gynecol* 2000;182:1033–5. [PubMed] ↩
48. Bariatric surgery and pregnancy. ACOG Practice Bulletin No. 105. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2009;113:1405–13. [PubMed] [Obstetrics & Gynecology] ↩
49. Wax JR, Wolff R, Cobean R, Pinette MG, Blackstone J, Cartin A. Intussusception complicating pregnancy following laparoscopic Roux-en-Y gastric bypass. *Obes Surg* 2007; 17:977–9. [PubMed] ↩
50. Patel JA, Patel NA, Thomas RL, Nelms JK, Colella JJ. Pregnancy outcomes after laparoscopic Roux-en-Y gastric bypass. *Surg Obes Relat Dis* 2008;4:39–45. [PubMed] ↩
51. Sheiner E, Levy A, Silverberg D, Menes TS, Levy I, Katz M, et al. Pregnancy after bariatric surgery is not associated with adverse perinatal outcome. *Am J Obstet Gynecol* 2004; 190:1335–40. [PubMed] [Full Text] ↩
52. Martin LF, Finigan KM, Nolan TE. Pregnancy after adjustable gastric banding. *Obstet Gynecol* 2000;95:927–30. [PubMed] [Obstetrics & Gynecology] ↩
53. Weiss HG, Nehoda H, Labeck B, Hourmont K, Marth C, Aigner F. Pregnancies after adjustable gastric banding. *Obes Surg* 2001;11:303–6. [PubMed] ↩
54. Maggard MA, Yermilov I, Li Z, Maglione M, Newberry S, Suttorp M, et al. Pregnancy and fertility following bariatric surgery: a systematic review. *JAMA* 2008;300:2286–96. [PubMed] [Full Text] ↩

Copyright January 2013 by the American College of Obstetricians and Gynecologists, 409 12th Street, SW, PO Box 96920, Washington, DC 20090-6920. All rights reserved.

ISSN 1074-861X

Obesity in pregnancy. Committee Opinion No. 549. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2013;121:213–7.